


CONFIDENTIAL LABORATORY REPORT

Member of Clinical and Laboratory Standards Institute, U.S.A.



Asiri Hospital Holdings PLC, 181, Kirula Road, Narahenpita, Colombo 05
T. +94 11 452 3355-7 F. +94 11 452 3358 priab@asiri.lk

**HISTOPATHOLOGY**

** OPD/AHH/ALS **

Page 2 of 2

REFERENCE No. : 01 0595 22/01/25
 SAMPLE DATE & TIME : 22/01/2025 18:41
 REPORT DATE & TIME : 31/01/2025 18:25 AGE : 52 Y/M
 PATIENT : MR. ROSHAN FERNANDO 104C UHID 310671227
 REFERRED BY : DR (MRS) LAKMALIE PARANAHEWA

10-20% of the surface areas of the affected cores from the left lobe. There is an extensive intraduct carcinoma component. There is no evidence of vascular invasion. Perineural invasion is present.

Conclusion :- A-F) TRUS guided prostatic biopsy.

- 1) Tumour type - Prostatic adenocarcinoma (acinar type)
- 2) Gleason sum score - $4 + 5 = 9$
- 3) Grade group (ISUP) - Group 5
- 4) Tumour burden -
 Percentage of positive biopsy cores - 75%.
 The tumour occupies 80-100% of the surface areas of the cores from the right lobe (average 93.3%) and 10-20% of the surface areas of the affected cores from the left lobe.
- 5) Vascular invasion - Absent
- 6) Perineural invasion - Present
- 7) Intraduct carcinoma component - Present.

PCH - 292

(S.C.T. 22/01/25 at 8.35 am)

Chandu de Silva

PROF. CHANDU DE SILVA
 MBBS, D.Path, MD (Histopathology), FCPPathSL, FCSSL, MIAC
 Professor of Pathology